
Landscape patterns of early postfire lodgepole pine regeneration dominate stand structure and function 24 years after the 1988 Yellowstone Fires

Monica G. Turner¹, William H. Romme², Daniel B. Tinker³, and Timothy G. Whitby¹
¹University of Wisconsin-Madison, ²Colorado State University, ³University of Wyoming





Background

- Understanding the rate and spatial variability of postfire succession in western US forests is important.
 - Increasing fire → more young, postfire forests
 - Few studies have tracked landscape patterns of succession following large, stand-replacing fires
- The 1988 Yellowstone Fires affected > 300,000 ha of forest, creating a complex postfire mosaic (Romme et al. 2011).
 - Striking landscape variation in Initial postfire lodgepole pine (*Pinus contorta*) regeneration, ranging from 0 to > 500,000 ha⁻¹ in response to prefire serotiny and fire severity (Turner 2010)

Question and hypotheses

- Are stand structure and function beginning to converge 24 years after the 1988 Fires?
 - H1: Landscape variation in postfire tree density and basal area have declined due to density-dependent mortality.
 - H2: Landscape variation in aboveground net primary production (ANPP) has decreased due to decreasing ANPP in high-density stands and increasing ANPP in low-to moderate-density stands.
 - Postfire tree density remains a "master variable" that determines stand-level ecosystem function.

Methods

• In 2012, we re-sampled 72 widely distributed 0.25-ha plots in postfire lodgepole pine last sampled in 1999 (11 years postfire), using the same protocols (Turner et al. 2004).






Results


H1 Stand structure: Tree density

- Landscape variation in postfire lodgepole pine density has changed little thus far.
 - Similar CV at 11 (285%) and 24 yrs (250%) postfire





 Lodgepole pine density changed relatively little from 11 to 24 yrs postfire and remained <u>very</u> high.




- Mean of 21,738 ha⁻¹ (range 0 to 344,000) at 24 yrs postfire
 - Decline in mean (from 32,000 ha⁻¹ at 11 yrs postfire) driven by very large declines (>100,000 ha⁻¹) in three highest-density stands
- Median increased by 233 ha⁻¹ since 11 yrs postfire
 - Tree density increased in 60% of the plots
 - Greater increase in low-density stands, but even stands with density >10,000 ha⁻¹ increased
- Lodgepole pine density at 24 yrs highly correlated with density at 11 yrs postfire (R = 0.90 P < 0.0001)

H1 Stand structure: Basal area


- Landscape variation in lodgepole pine basal area also remained very high at 24 yrs postfire.
 - Mean of 25.3 m² ha⁻¹ (range 0 to146), increasing with stand density (R = 0.87, P < 0.0001)
 - Dense stands have smaller trees
 (R = -0.71, P < 0.0001)






H2 Stand function: ANPP


- Landscape variation in ANPP declined between 11 yrs (CV 156%) and 24 yrs (CV 106%) postfire.
- Mean lodgepole pine ANPP increased ~10x between 11 and 24 yrs postfire (from 1.72 to 11.92 Mg ha⁻¹ yr⁻¹).
 - ANPP increased in all stands with trees
 - ANPP increased with tree density (R = 0.76, P < 0.0001)





- Mean herbaceous ANPP increased ~40% between 11 and 24 yrs postfire (from 1.05 to 1.44 Mg ha⁻¹ yr⁻¹).
 - Herbaceous ANPP declined with tree density





Conclusions

- Stand structure is not yet converging, but ANPP is increasing rapidly and shows signs of converging.
- Initial landscape pattern of postfire tree density still dominates stand trajectories 25 yrs after the '88 fires.
 - Contingent factors → persistent landscape legacy





Literature cited

Romme, W. H., M. S. Boyce, R. E. Gresswell, E. H. Merrill, G. W. Minshall, C. Whitlock and M. G. Turner. 2011. Twenty years after the 1988 Yellowstone fires: lessons about disturbance and ecosystems. Ecosystems 14:1196-1215.

Turner, M. G. 2010. Disturbance and landscape dynamics in a changing world. Ecology 91:2833-2849.

Turner, M. G., D. B. Tinker, W. H. Romme, D. M. Kashian and C. M. Litton. 2004. Landscape patterns of sapling density, leaf area, and aboveground net primary production in postfire lodgepole pine forests, Yellowstone National Park (USA). Ecosystems 7:751-775.

Acknowledgements

 Thanks to Kellen Nelson, Paige Copenhaver, Dan Donato, Natalie Kaner, Andy Muench and Winslow Hansen for field assistance during summer 2012.



Research funded by the Joint Fire Science Program